

**Table 1.** Clinical characteristics according to H<sub>2</sub>FPEF score by BMI  $\geq$  30 kg/m<sup>2</sup>

		<b>H<sub>2</sub>FPEF score</b>			<b>P value</b>	
		<b>All patients (n = 497)</b>	<b>Low-score group (0-3 point) (n = 167)</b>	<b>Intermediate-score group (4-6 point) (n = 230)</b>		<b>High-score group (7-9 point) (n = 100)</b>
<b>H<sub>2</sub>FPEF score</b>						
<b>Heavy</b>	(2pts)	79 (15.9)	8 (4.8)	28 (12.2)	43 (43.0)	<0.001
<b>BMI <math>\geq</math> 30 kg/m<sup>2</sup> (%)</b>						
<b>Hypertension</b>	(1pt)	299 (60.2)	81 (48.5)	144 (62.6)	88 (88.0)	<0.001
<b><math>\geq</math> 2 antihypertensive drugs (%)</b>						
<b>Atrial Fibrillation</b>	(3pts)	313 (63.0)	5 (3.0)	166 (72.2)	100 (100.0)	<0.001
<b>History of AF (%)</b>						
<b>Pulmonary hypertension</b>	(1pt)	271 (54.5)	25 (15.0)	84 (36.5)	72 (72.0)	<0.001
<b>PAP <math>\geq</math> 35 mmHg (%)</b>						
<b>Elder</b>	(1pt)	181 (36.4)	126 (75.4)	190 (82.6)	93 (93.0)	<0.001
<b>Age <math>\geq</math> 60 years (%)</b>						
<b>Filling Pressure</b>	(1pt)	409 (82.3)	12 (77.2)	204 (88.7)	96 (96.0)	<0.001
<b>E/e' (septal) <math>\geq</math> 9 (%)</b>						

Data are number of patients (%). HFpEF: heart failure with preserved ejection fraction; BMI: body mass index; AF: paroxysmal/persistent atrial fibrillation; PAP: pulmonary artery systolic pressure; E/e' (septal): the ratio of early transmitral flow velocity to tissue Doppler early diastolic mitral annular velocity (septal).